



Ampicillin Reconstitution Volumes




| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes




| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes




| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes




| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes




| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes




| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes



| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |

Ampicillin Reconstitution Volumes



| | | |
|---------|--------|-------------|
| 250 mg | -----> | 2.5 ml SWFI |
| 500 mg | -----> | 5 ml SWFI |
| 1000 mg | -----> | 10 ml SWFI |
| 2000 mg | -----> | 20 ml SWFI |